

Charmonium mass in hot asymmetric nuclear matter

Amruta Mishra* and Arvind Kumar†

*Department of Physics, Indian Institute of Technology,
Delhi, Hauz Khas, New Delhi – 110 016, India*

Abstract

We calculate the in-medium masses of J/ψ and of the excited states of charmonium ($\psi(3686)$ and $\psi(3770)$) in isospin asymmetric nuclear matter at finite temperatures. These mass modifications arise due to the interaction of the charmonium states with the gluon condensates of QCD, simulated by a scalar dilaton field introduced to incorporate the broken scale invariance of QCD within an effective chiral model. The change in the mass of J/ψ in the nuclear matter with density is seen to be rather small, as has been shown in the literature by using various approaches, whereas, the masses of the excited states of charmonium ($\psi(3686)$ and $\psi(3770)$) are seen to have considerable drop at high densities. The dependence of the masses of the charmonium states on the isospin asymmetry has also been investigated in the hot nuclear matter and is seen to be appreciable at moderate temperatures and high densities. These medium modifications of the charmonium states should modify the experimental observables arising from the compressed baryonic matter produced in asymmetric heavy ion collision experiments in the future facility of FAIR, GSI.

PACS numbers: 24.10.Cn; 13.75.Jz; 25.75.-q

*Electronic address: amruta@physics.iitd.ac.in, mishra@th.physik.uni-frankfurt.de

†Electronic address: iitd.arvind@gmail.com

I. INTRODUCTION

The study of the in-medium properties of hadrons is an active topic of research in strong interaction physics, both experimentally and theoretically. The topic is of direct relevance in the context of heavy ion collision experiments, which probe matter at high temperatures and/or densities. The medium modifications of the hadrons have direct consequences on the experimental observables from the strongly interacting matter produced in heavy ion collision experiments. The in-medium properties of the kaons and antikaons in the nuclear medium are of relevance in neutron star phenomenology where an attractive interaction of antikaon-nucleon can lead to antikaon condensation in the interior of the neutron stars. The medium modifications of kaons and antikaons also show in their production and propagation in the heavy ion collision experiments. The modifications of the properties of the charm mesons, D and $\bar{D}$ as well as the J/ψ mesons and the excited states of charmonium, can have important consequences on the production of open charm and the suppression of J/ψ in heavy ion collision experiments. In high energy heavy ion collision experiments at RHIC as well as LHC, the suppression of J/ψ can arise from the formation of the quark-gluon-plasma (QGP) [1, 2].

The D ($\bar{D}$) mesons are made up of one heavy charm quark (antiquark) and one light (u or d) antiquark (quark). In the QCD sum rule calculations, the mass modifications of D ($\bar{D}$) mesons in the nuclear medium arise due to the interactions of light antiquark (quark) present in the $D(\bar{D})$ mesons with the light quark condensate. There is appreciable change in the light quark condensate in the nuclear medium and hence D ($\bar{D}$) meson mass, due to its interaction with the light quark condensate, changes appreciably in the hadronic matter. On the other hand, the charmonium states are made up of a heavy charm quark and a charm antiquark. Within QCD sum rules, it is suggested that these heavy charmonium states interact with the nuclear medium through the gluon condensates. This is contrary to the interaction of the light vector mesons (ρ , ω , ϕ), which interact with the nuclear medium through the quark condensates. This is because all the heavy quark condensates can be related to the gluon condensates via heavy-quark expansion [3]. Also in the nuclear medium there are no valence charm quarks to leading order in density and any interaction with the medium is gluonic. The QCD sum rules has been used to study the medium modification of

D mesons [4] and light vector mesons [5]. The QCD sum rule approach [6] and leading order perturbative calculations [7] to study the medium modifications of charmonium, show that the mass of J/ψ is reduced only slightly in the nuclear medium. In [8], the mass modification of charmonium has been studied using leading order QCD formula and the linear density approximation for the gluon condensate in the nuclear medium. This shows a small drop for the J/ψ mass at the nuclear matter density, but there is seen to be significant shift in the masses of the excited states of charmonium ($\psi(3686)$ and $\psi(3770)$).

In the present work, we study the medium modification of the masses of J/ψ and excited charmonium states $\psi(3686)$ and $\psi(3770)$ in the nuclear medium due to the interaction with the gluon condensates using the leading order QCD formula. The gluon condensate in the nuclear medium is calculated from the medium modification of a scalar dilaton field introduced within a chiral $SU(3)$ model [9] through a scale symmetry breaking term in the Lagrangian density leading to the QCD trace anomaly. In the chiral $SU(3)$ model, the gluon condensate is related to the fourth power of the dilaton field χ and the changes in the dilaton field with density are seen to be small. The model has been used successfully to study the medium modifications of kaons and antikaons in isospin asymmetric nuclear matter in [10] and in hyperonic matter in [11]. The model has also been used to study the D mesons in asymmetric nuclear matter at zero temperature [12] and in the symmetric and asymmetric nuclear matter at finite temperatures in Ref.[13] and [14]. The vector mesons have also been studied within the model [15, 16]. In the present investigation, we study the isospin dependence of the in-medium masses of charmonium obtained from the dilaton field, χ calculated for the asymmetric nuclear matter at finite temperatures. This study will be of relevance for the experimental observables from high density matter produced in the asymmetric nuclear collisions at the future facility at GSI.

The outline of the paper is as follows : In section II, we give a brief introduction of the chiral $SU(3)$ model used to study the in-medium masses of charmonium in the present investigation. The medium modifications of the charmonium masses arise from the medium modification of a scalar dilaton field introduced in the hadronic model to incorporate broken scale invariance of QCD leading to QCD trace anomaly. In section III, we summarize the results obtained in the present investigation.

II. THE HADRONIC CHIRAL $SU(3) \times SU(3)$ MODEL

We use an effective chiral $SU(3)$ model for the present investigation [9]. The model is based on the nonlinear realization of chiral symmetry [17–19] and broken scale invariance [9, 15, 16]. This model has been used successfully to describe nuclear matter, finite nuclei, hypernuclei and neutron stars. The effective hadronic chiral Lagrangian density contains the following terms

$$\mathcal{L} = \mathcal{L}_{kin} + \sum_{W=X,Y,V,A,u} \mathcal{L}_{BW} + \mathcal{L}_{vec} + \mathcal{L}_0 + \mathcal{L}_{SB} \quad (1)$$

In Eq. (1), $\mathcal{L}_{kin}$ is kinetic energy term, $\mathcal{L}_{BW}$ is the baryon-meson interaction term in which the baryon-spin-0 meson interaction term generates the vacuum baryon masses. $\mathcal{L}_{vec}$ describes the dynamical mass generation of the vector mesons via couplings to the scalar mesons and contain additionally quartic self-interactions of the vector fields. $\mathcal{L}_0$ contains the meson-meson interaction terms inducing the spontaneous breaking of chiral symmetry as well as a scale invariance breaking logarithmic potential. $\mathcal{L}_{SB}$ describes the explicit chiral symmetry breaking.

To study the hadron properties at finite temperature and densities in the present investigation, we use the mean field approximation, where all the meson fields are treated as classical fields. In this approximation, only the scalar and the vector fields contribute to the baryon-meson interaction, $\mathcal{L}_{BW}$ since for all the other mesons, the expectation values are zero. The interactions of the scalar mesons and vector mesons with the baryons are given as

$$\mathcal{L}_{Bscal} + \mathcal{L}_{Bvec} = - \sum_i \bar{\psi}_i [m_i^* + g_{\omega i} \gamma_0 \omega + g_{\rho i} \gamma_0 \rho + g_{\phi i} \gamma_0 \phi] \psi_i. \quad (2)$$

The interaction of the vector mesons, of the scalar fields and the interaction corresponding to the explicitly symmetry breaking in the mean field approximation are given as

$$\begin{aligned} \mathcal{L}_{vec} = & \frac{1}{2} \left(m_\omega^2 \omega^2 + m_\rho^2 \rho^2 + m_\phi^2 \phi^2 \right) \frac{\chi^2}{\chi_0^2} \\ & + g_4 (\omega^4 + 6\omega^2 \rho^2 + \rho^4 + 2\phi^4), \end{aligned} \quad (3)$$

$$\mathcal{L}_0 = -\frac{1}{2} k_0 \chi^2 (\sigma^2 + \zeta^2 + \delta^2) + k_1 (\sigma^2 + \zeta^2 + \delta^2)^2$$

$$\begin{aligned}
& + k_2 \left(\frac{\sigma^4}{2} + \frac{\delta^4}{2} + 3\sigma^2\delta^2 + \zeta^4 \right) + k_3\chi \left(\sigma^2 - \delta^2 \right) \zeta \\
& - k_4\chi^4 - \frac{1}{4}\chi^4 \ln \frac{\chi^4}{\chi_0^4} + \frac{d}{3}\chi^4 \ln \left(\left(\frac{(\sigma^2 - \delta^2)\zeta}{\sigma_0^2\zeta_0} \right) \left(\frac{\chi}{\chi_0} \right)^3 \right),
\end{aligned} \tag{4}$$

and

$$\mathcal{L}_{SB} = - \left(\frac{\chi}{\chi_0} \right)^2 \left[m_\pi^2 f_\pi \sigma + \left(\sqrt{2} m_k^2 f_k - \frac{1}{\sqrt{2}} m_\pi^2 f_\pi \right) \zeta \right]. \tag{5}$$

The effective mass of the baryon of species i is given as

$$m_i^* = -(g_{\sigma i} \sigma + g_{\zeta i} \zeta + g_{\delta i} \delta) \tag{6}$$

The baryon-scalar meson interactions, as can be seen from equation (6), generate the baryon masses through the coupling of baryons to the non-strange σ , strange ζ scalar mesons and also to scalar-isovector meson δ . In analogy to the baryon-scalar meson coupling there exist two independent baryon-vector meson interaction terms corresponding to the F-type (antisymmetric) and D-type (symmetric) couplings. Here antisymmetric coupling is used because the universality principle [20] and vector meson dominance model suggest small symmetric coupling. Additionally, we choose the parameters [9, 10] so as to decouple the strange vector field $\phi_\mu \sim \bar{s} \gamma_\mu s$ from the nucleon, corresponding to an ideal mixing between ω and ϕ mesons. A small deviation of the mixing angle from ideal mixing [21–23] has not been taken into account in the present investigation.

The concept of broken scale invariance leading to the trace anomaly in (massless) QCD, $\theta_\mu^\mu = \frac{\beta_{QCD}}{2g} G_{\mu\nu}^a G^{\mu\nu a}$, where $G_{\mu\nu}^a$ is the gluon field strength tensor of QCD, is simulated in the effective Lagrangian at tree level [24] through the introduction of the scale breaking terms

$$\mathcal{L}_{scalebreaking} = -\frac{1}{4}\chi^4 \ln \left(\frac{\chi^4}{\chi_0^4} \right) + \frac{d}{3}\chi^4 \ln \left(\left(\frac{I_3}{\det \langle X \rangle_0} \right) \left(\frac{\chi}{\chi_0} \right)^3 \right), \tag{7}$$

where $I_3 = \det \langle X \rangle$, with X as the multiplet for the scalar mesons. These scale breaking terms, in the mean field approximation, are given by the last two terms of the Lagrangian density, $\mathcal{L}_0$ given by equation (4). The effect of these logarithmic terms is to break the scale invariance, which leads to the trace of the energy momentum tensor as [25]

$$\theta_\mu^\mu = \chi \frac{\partial \mathcal{L}}{\partial \chi} - 4\mathcal{L} = -(1-d)\chi^4. \tag{8}$$

Hence the scalar gluon condensate of QCD ($\langle G_{\mu\nu}^a G^{\mu\nu a} \rangle$) is simulated by a scalar dilaton field in the present hadronic model.

The coupled equations of motion for the non-strange scalar field σ , strange scalar field ζ , scalar-isovector field δ and dilaton field χ , are derived from the Lagrangian density and are given as

$$k_0\chi^2\sigma - 4k_1(\sigma^2 + \zeta^2 + \delta^2)\sigma - 2k_2(\sigma^3 + 3\sigma\delta^2) - 2k_3\chi\sigma\zeta - \frac{d}{3}\chi^4\left(\frac{2\sigma}{\sigma^2 - \delta^2}\right) + \left(\frac{\chi}{\chi_0}\right)^2 m_\pi^2 f_\pi - \sum g_{\sigma i} \rho_i^s = 0 \quad (9)$$

$$k_0\chi^2\zeta - 4k_1(\sigma^2 + \zeta^2 + \delta^2)\zeta - 4k_2\zeta^3 - k_3\chi(\sigma^2 - \delta^2) - \frac{d}{3}\frac{\chi^4}{\zeta} + \left(\frac{\chi}{\chi_0}\right)^2 \left[\sqrt{2}m_k^2 f_k - \frac{1}{\sqrt{2}}m_\pi^2 f_\pi \right] - \sum g_{\zeta i} \rho_i^s = 0 \quad (10)$$

$$k_0\chi^2\delta - 4k_1(\sigma^2 + \zeta^2 + \delta^2)\delta - 2k_2(\delta^3 + 3\sigma^2\delta) + k_3\chi\delta\zeta + \frac{2}{3}d\left(\frac{\delta}{\sigma^2 - \delta^2}\right) - \sum g_{\delta i} \rho_i^s = 0 \quad (11)$$

$$k_0\chi(\sigma^2 + \zeta^2 + \delta^2) - k_3(\sigma^2 - \delta^2)\zeta + \chi^3 \left[1 + \ln\left(\frac{\chi^4}{\chi_0^4}\right) \right] + (4k_4 - d)\chi^3 - \frac{4}{3}d\chi^3 \ln\left(\left(\frac{(\sigma^2 - \delta^2)\zeta}{\sigma_0^2\zeta_0}\right)\left(\frac{\chi}{\chi_0}\right)^3\right) + \frac{2\chi}{\chi_0^2} \left[m_\pi^2 f_\pi \sigma + \left(\sqrt{2}m_k^2 f_k - \frac{1}{\sqrt{2}}m_\pi^2 f_\pi \right) \zeta \right] = 0 \quad (12)$$

In the above, ρ_i^s are the scalar densities for the baryons, given as

$$\rho_i^s = \gamma_i \int \frac{d^3k}{(2\pi)^3} \frac{m_i^*}{E_i^*(k)} \left(\frac{1}{e^{(E_i^*(k) - \mu_i^*)/T} + 1} + \frac{1}{e^{(E_i^*(k) + \mu_i^*)/T} + 1} \right) \quad (13)$$

where, $E_i^*(k) = (k^2 + m_i^{*2})^{1/2}$, and, $\mu_i^* = \mu_i - g_{\omega i}\omega - g_{\rho i}\rho - g_{\phi i}\phi$, are the single particle energy and the effective chemical potential for the baryon of species i , and, $\gamma_i=2$ is the spin degeneracy factor [10].

The above coupled equations of motion are solved to obtain the density and temperature dependent values of the scalar fields (σ , ζ and δ) and the dilaton field, χ , in the isospin asymmetric hot nuclear medium. As has been already mentioned, the value of the χ is related to the scalar gluon condensate in the hot hadronic medium, and is used to compute the in-medium masses of charmonium states, in the present investigation. The isospin asymmetry

in the medium is introduced through the scalar-isovector field δ and therefore the dilaton field obtained after solving the above equations is also dependent on the isospin asymmetry parameter, η defined as $\eta = (\rho_n - \rho_p)/(2\rho_B)$, where ρ_n and ρ_p are the number densities of the neutron and the proton and ρ_B is the baryon density. In the present investigation, we study the effect of isospin asymmetry of the medium on the masses of the charmonium states J/ψ , $\psi(3686)$ and $\psi(3770)$.

The comparison of the trace of the energy momentum tensor arising from the trace anomaly of QCD with that of the present chiral model gives the relation of the dilaton field to the scalar gluon condensate. We have, in the limit of massless quarks [26],

$$\theta_\mu^\mu = \langle \frac{\beta_{QCD}}{2g} G_{\mu\nu}^a G^{\mu\nu a} \rangle \equiv -(1-d)\chi^4 \quad (14)$$

The parameter d originates from the second logarithmic term of equation (7). To get an insight into the value of the parameter d , we recall that the QCD β function at one loop level, for N_c colors and N_f flavors is given by

$$\beta_{QCD}(g) = -\frac{11N_c g^3}{48\pi^2} \left(1 - \frac{2N_f}{11N_c}\right) + O(g^5) \quad (15)$$

In the above equation, the first term in the parentheses arises from the (antiscreening) self-interaction of the gluons and the second term, proportional to N_f , arises from the (screening) contribution of quark pairs. Equations (14) and (15) suggest the value of d to be $6/33$ for three flavors and three colors, and for the case of three colors and two flavors, the value of d turns out to be $4/33$, to be consistent with the one loop estimate of QCD β function. These values give the order of magnitude about which the parameter d can be taken [25], since one cannot rely on the one-loop estimate for $\beta_{QCD}(g)$. In the present investigation of the in-medium properties of the charmonium states due to the medium modification of the dilaton field within chiral $SU(3)$ model, we use the value of $d=0.064$ [12]. This parameter, along with the other parameters corresponding to the scalar Lagrangian density, $\mathcal{L}_0$ given by (4), are fitted so as to ensure extrema in the vacuum for the σ , ζ and χ field equations, to reproduce the vacuum masses of the η and η' mesons, the mass of the σ meson around 500 MeV, and, pressure, $p(\rho_0)=0$, with ρ_0 as the nuclear matter saturation density [9, 12].

The trace of the energy-momentum tensor in QCD, using the one loop beta function

given by equation (15), for $N_c=3$ and $N_f=3$, is given as,

$$\theta_\mu^\mu = -\frac{9}{8} \frac{\alpha_s}{\pi} G_{\mu\nu}^a G^{\mu\nu a} \quad (16)$$

Using equations (14) and (16), we can write

$$\left\langle \frac{\alpha_s}{\pi} G_{\mu\nu}^a G^{\mu\nu a} \right\rangle = \frac{8}{9} (1-d) \chi^4 \quad (17)$$

We thus see from the equation (17) that the scalar gluon condensate $\left\langle \frac{\alpha_s}{\pi} G_{\mu\nu}^a G^{\mu\nu a} \right\rangle$ is proportional to the fourth power of the dilaton field, χ , in the chiral SU(3) model. As mentioned earlier, the in-medium masses of charmonium states are modified due to the gluon condensates. Therefore, we need to know the change in the gluon condensate with density and temperature of the asymmetric nuclear medium, which is calculated from the modification of the χ field, by using equation (17).

From the QCD sum rule calculations, the mass shift of the charmonium states in the medium is due to the gluon condensates [4, 8]. For heavy quark systems, there are two independent lowest dimension operators: the scalar gluon condensate ($\left\langle \frac{\alpha_s}{\pi} G_{\mu\nu}^a G^{\mu\nu a} \right\rangle$) and the condensate of the twist 2 gluon operator ($\left\langle \frac{\alpha_s}{\pi} G_{\mu\nu}^a G^{\mu\alpha a} \right\rangle$). These operators can be rewritten in terms of the color electric and color magnetic fields, $\left\langle \frac{\alpha_s}{\pi} \vec{E}^2 \right\rangle$ and $\left\langle \frac{\alpha_s}{\pi} \vec{B}^2 \right\rangle$. Additionally, since the Wilson coefficients for the operator $\left\langle \frac{\alpha_s}{\pi} \vec{B}^2 \right\rangle$ vanishes in the non-relativistic limit, the only contribution from the gluon condensates is proportional to $\left\langle \frac{\alpha_s}{\pi} \vec{E}^2 \right\rangle$, similar to the second order Stark effect. Hence, the mass shift of the charmonium states arises due to the change in the operator $\left\langle \frac{\alpha_s}{\pi} \vec{E}^2 \right\rangle$ in the medium from its vacuum value [8]. In the leading order mass shift formula derived in the large charm mass limit [7], the shift in the mass of the charmonium state is given as [8]

$$\Delta m_\psi(\epsilon) = -\frac{1}{9} \int dk^2 \left| \frac{\partial \psi(k)}{\partial k} \right|^2 \frac{k}{k^2/m_c + \epsilon} \left(\left\langle \frac{\alpha_s}{\pi} E^2 \right\rangle - \left\langle \frac{\alpha_s}{\pi} E^2 \right\rangle_0 \right). \quad (18)$$

In the above, m_c is the mass of the charm quark, taken as 1.95 GeV [8], m_ψ is the vacuum mass of the charmonium state and $\epsilon = 2m_c - m_\psi$. $\psi(k)$ is the wave function of the charmonium state in the momentum space, normalized as $\int \frac{d^3k}{2\pi^3} |\psi(k)|^2 = 1$ [27]. At finite densities, in the linear density approximation, the change in the value of $\left\langle \frac{\alpha_s}{\pi} \vec{E}^2 \right\rangle$, from its vacuum value, is given as

$$\left\langle \frac{\alpha_s}{\pi} E^2 \right\rangle - \left\langle \frac{\alpha_s}{\pi} E^2 \right\rangle_0 = \left\langle \frac{\alpha_s}{\pi} E^2 \right\rangle_N \frac{\rho_B}{2M_N}, \quad (19)$$

and the mass shift in the charmonium states reduces to [8]

$$\Delta m_\psi(\epsilon) = -\frac{1}{9} \int dk^2 \left| \frac{\partial \psi(k)}{\partial k} \right|^2 \frac{k}{k^2/m_c + \epsilon} \left\langle \frac{\alpha_s}{\pi} E^2 \right\rangle_N \frac{\rho_B}{2M_N}. \quad (20)$$

In the above, $\left\langle \frac{\alpha_s}{\pi} E^2 \right\rangle_N$ is the expectation value of $\left\langle \frac{\alpha_s}{\pi} E^2 \right\rangle$ with respect to the nucleon.

The expectation value of the scalar gluon condensate can be expressed in terms of the color electric field and the color magnetic field as [28]

$$\left\langle \frac{\alpha_s}{\pi} G_{\mu\nu}^a G^{\mu\nu a} \right\rangle = -2 \left\langle \frac{\alpha_s}{\pi} (E^2 - B^2) \right\rangle. \quad (21)$$

In the non-relativistic limit, as already mentioned, the contribution from the magnetic field vanishes and hence, we can write,

$$\left\langle \frac{\alpha_s}{\pi} E^2 \right\rangle = -\frac{1}{2} \left\langle \frac{\alpha_s}{\pi} G_{\mu\nu}^a G^{\mu\nu a} \right\rangle \quad (22)$$

Using equations (17), (18) and (22), we obtain the expression for the mass shift in the charmonium in the hot and dense nuclear medium, which arises from the change in the dilaton field in the present investigation, as

$$\Delta m_\psi(\epsilon) = \frac{4}{81} (1-d) \int dk^2 \left| \frac{\partial \psi(k)}{\partial k} \right|^2 \frac{k}{k^2/m_c + \epsilon} (\chi^4 - \chi_0^4). \quad (23)$$

In the above, χ and χ_0 are the values of the dilaton field in the nuclear medium and the vacuum respectively.

In the present investigation, the wave functions for the charmonium states are taken to be Gaussian and are given as [29]

$$\psi_{N,l} = \text{Normalization} \times Y_l^m(\theta, \phi) (\beta^2 r^2)^{\frac{1}{2}l} \exp^{-\frac{1}{2}\beta^2 r^2} L_{N-1}^{l+\frac{1}{2}}(\beta^2 r^2) \quad (24)$$

where $\beta^2 = M\omega/h$ characterizes the strength of the harmonic potential, $M = m_c/2$ is the reduced mass of the charm quark and charm anti-quark system, and $L_p^k(z)$ is the associated Laguerre Polynomial. As in Ref. [8], the oscillator constant β is determined from the mean squared radii $\langle r^2 \rangle$ as 0.46^2 fm^2 , 0.96^2 fm^2 and 1 fm^2 for the charmonium states $J/\psi(3097)$, $\psi(3686)$ and $\psi(3770)$, respectively. This gives the value for the parameter β as 0.51 GeV , 0.38 GeV and 0.37 GeV for $J/\psi(3097)$, $\psi(3686)$ and $\psi(3770)$, assuming that these charmonium states are in the 1S, 2S and 1D states respectively. Knowing the wave functions

of the charmonium states and calculating the medium modification of the dilaton field in the hot nuclear matter, we obtain the mass shift of the charmonium states, J/ψ , $\psi(3686)$ and $\psi(3770)$ respectively. In the next section we shall present the results of the present investigation of these in-medium charmonium masses in hot asymmetric nuclear matter.

III. RESULTS AND DISCUSSIONS

In this section, we first investigate the effects of density, isospin-asymmetry and temperature of the nuclear medium on the dilaton field χ using a chiral SU(3) model. From the medium modification of the χ field, we shall then study the in-medium masses of charmonium states J/ψ , $\psi(3686)$ and $\psi(3770)$ using equation (23).

The values of the parameters used in the present investigation, are : $k_0 = 2.54$, $k_1 = 1.35$, $k_2 = -4.78$, $k_3 = -2.77$, $k_4 = -0.22$ and $d = 0.064$, which are the parameters occurring in the scalar meson interactions defined in equation (4). The vacuum values of the scalar isoscalar fields, σ and ζ and the dilaton field χ are -93.3 MeV, -106.6 MeV and 409.77 MeV respectively. The values, $g_{\sigma N} = 10.6$ and $g_{\zeta N} = -0.47$ are determined by fitting to the vacuum baryon masses. The other parameters fitted to the asymmetric nuclear matter saturation properties in the mean-field approximation are: $g_{\omega N} = 13.3$, $g_{\rho p} = 5.5$, $g_4 = 79.7$, $g_{\delta p} = 2.5$, $m_\zeta = 1024.5$ MeV, $m_\sigma = 466.5$ MeV and $m_\delta = 899.5$ MeV. The nuclear matter saturation density used in the present investigation is 0.15 fm^{-3} .

The variations of the dilaton field χ with density, temperature and isospin asymmetry, within the chiral SU(3) model, are obtained by solving the coupled equations of motion of scalar fields given by equations (9), (10), (11) and (12). In figure 1, we show the variation of dilaton field χ , with temperature, for both zero and finite baryon densities, and for selected values of the isospin asymmetry parameter, $\eta = 0, 0.1, 0.3$ and 0.5 . At zero baryon density, it is observed that the value of the dilaton field remains almost a constant upto a temperature of about 130 MeV above which it is seen to drop with increase in temperature. However, the drop in the dilaton field is seen to be very small. The value of the dilaton field is seen to change from 409.8 MeV at $T=0$ to about 409.7 MeV and 409.3 MeV at $T=150$ MeV and $T=175$ MeV respectively. The thermal distribution functions have an effect of increasing the scalar densities at zero baryon density, i.e., for $\mu_i^*=0$, as can be seen from the expression




FIG. 1: (Color online) The dilaton field χ plotted as a function of the temperature, at given baryon densities, for different values of the isospin asymmetry parameter, η .

of the scalar densities, given by (13). This effect seems to be negligible up to a temperature of about 130 MeV. This leads to a decrease in the magnitudes of scalar fields, σ and ζ . This behaviour of the scalar fields is reflected in the value of χ , which is solved from the coupled equations of motion of the scalar fields, given by equations (9), (10), (11) and (12), as a drop as we increase the temperature above a temperature of about 130 MeV. The scalar densities

attaining nonzero values at high temperatures, even at zero baryon density, indicates the presence of baryon-antibaryon pairs in the thermal bath and has already been observed in the literature [16, 30]. This leads to the baryon masses to be different from their vacuum masses above this temperature, arising from modifications of the scalar fields σ and ζ .

For finite density situations, the behaviour of the χ field with temperature is seen to be very different from the zero density case, as can be seen from the subplots (b),(c) and (d) of figure 1, where the χ field is plotted as a function of the temperature for densities ρ_0 , $2\rho_0$ and $4\rho_0$ respectively. At finite densities, one observes first a rise and then a decrease of the dilaton field with temperature. This is related to the fact that at finite densities, the magnitude of the σ field (as well as of the ζ field) first show an increase and then a drop with further increase of the temperature [14] which is reflected in the behaviour of χ field, since it is solved from the coupled equations of the scalar fields. The reason for the different behaviour of the scalar fields (σ and ζ) at zero and finite densities can be understood in the following manner [16]. As has been already mentioned, the thermal distribution functions in (13) have an effect of increasing the scalar densities at zero baryon density, i.e., for $\mu_i^*=0$. However, at finite densities, i.e., for nonzero values of the effective chemical potential, μ_i^* , for increasing temperature, there are contributions also from higher momenta, thereby, increasing the denominator of the integrand on the right hand side of the equation (13). This leads to a decrease in the scalar density. The competing effects of the thermal distribution functions and the contributions of the higher momenta states give rise to the observed effect of the scalar density and hence of the σ and ζ fields with temperature at finite baryon densities [16]. This kind of behaviour of the scalar σ field on temperature at finite densities has also been observed in the Walecka model by Li and Ko [31], which was reflected as an increase in the mass of the nucleon with temperature at finite densities in the mean field calculations. The effects of the behaviour of the scalar fields on the value of the χ field, obtained from solving the coupled equations (9) to (12) for the scalar fields, are shown in figure 1.

In figure 1, it is observed that for a given value of isospin asymmetry parameter η , the dilaton field χ decreases with increase in the density of the nuclear medium. The drop in the value of χ with density is seen to be much larger as compared to its modification

with temperature at a given density. For isospin symmetric nuclear medium ($\eta = 0$) at temperature $T = 0$, the reduction in the dilaton field χ from its vacuum value ($\chi_0 = 409.8$ MeV), is seen to be about 3 MeV at $\rho_B = \rho_0$ and about 13 MeV, for $\rho_B = 4\rho_0$. As we move from isospin symmetric medium, with $\eta = 0$, to isospin asymmetric medium, at temperature $T = 0$, and, for a given value of density, there is seen to be an increase in the value of the dilaton field χ . However, the effect of isospin asymmetry of the medium on the value of the dilaton field is observed to be negligible upto about a density of nuclear matter saturation density, and is appreciable only at higher values of densities as can be seen in figure 1. At nuclear matter saturation density, ρ_0 , the value of dilaton field χ changes from 406.4 MeV in symmetric nuclear medium ($\eta = 0$) to 406.5 MeV in the isospin asymmetric nuclear medium ($\eta = 0.5$). At a density of about $4\rho_0$, the values of the dilaton field are modified to 396.7 MeV and 398 MeV at $\eta = 0$ and 0.5, respectively. Thus the increase in the dilaton field χ with isospin asymmetry of the medium is seen to be more at zero temperature as we move to higher densities.

At a finite density, ρ_B , and for given isospin asymmetry parameter η , the dilaton field χ is seen to first increase with temperature and above a particular value of the temperature, it is seen to decrease with further increase in temperature. At the nuclear saturation density $\rho_B = \rho_0$ and in isospin symmetric nuclear medium ($\eta = 0$) the value of the dilaton field χ increases upto a temperature of about $T = 145$ MeV, above with there is a drop in the dilaton field. For $\rho_B = \rho_0$ in the asymmetric nuclear matter with $\eta = 0.5$, there is seen to be a rise in the value of χ upto a temperature of about 120 MeV, above which it starts decreasing. As it has been already mentioned, at zero temperature and for a given value of density, the dilaton field χ is found to increase with increase in the isospin asymmetry of the nuclear medium. But from figure 1, it is observed that at high temperatures and for a given density, the value of the dilaton field χ becomes higher in symmetric nuclear medium as compared to isospin asymmetric nuclear medium e.g. at nuclear saturation density $\rho_B = \rho_0$ and temperature $T = 150$ MeV the values of dilaton field χ are 407.3 MeV and 407 MeV at $\eta = 0$ and 0.5 respectively. At density $\rho_B = 4\rho_0$, $T = 150$ MeV the values of dilaton field χ are seen to be 399.1 MeV and 398.7 MeV for $\eta = 0$ and 0.5 respectively. This observed behaviour of the χ is related to the fact that at finite densities and for isospin asymmetric




FIG. 2: (Color online) The mass shift of J/ψ plotted as a function of the baryon density in units of nuclear matter saturation density at given temperatures, for different values of the isospin asymmetry parameter, η .

matter, there are contributions from the scalar isovector δ field, whose magnitude is seen to decrease for higher temperatures for given densities, whereas δ field has zero contribution for isospin symmetric matter.

We shall now investigate how the behaviour of the dilaton field χ in the hot asymmetric




FIG. 3: (Color online) The mass shift of $\psi(3686)$ plotted as a function of the baryon density in units of nuclear matter saturation density at given temperatures, for different values of the isospin asymmetry parameter, η .

nuclear medium affects the in-medium masses of the charmonium states J/ψ , $\psi(3686)$ and $\psi(3770)$. In figures 2, 3 and 4, we show the shifts of the masses of charmonium states J/ψ , $\psi(3686)$ and $\psi(3770)$ from their vacuum values, as functions of the baryon density for given values of temperature T , and for different values of the isospin asymmetry parameter, η .




FIG. 4: (Color online) The mass shift of $\psi(3770)$ plotted as a function of the baryon density in units of nuclear matter saturation density at given temperatures, for different values of the isospin asymmetry parameter, η .

We have shown the results for the values of the temperature, $T = 0, 50, 100$ and 150 MeV. At the nuclear matter saturation density, $\rho_B = \rho_0$ at temperature $T = 0$, the mass-shift for J/ψ meson is observed to be about -8.6 MeV in the isospin symmetric nuclear medium ($\eta = 0$) and in the asymmetric nuclear medium, with isospin asymmetry parameter $\eta = 0.5$,

it is seen to be -8.4 MeV. At $\rho_B = 4\rho_0$, temperature $T = 0$, the mass-shift for J/ψ meson is -32.2 MeV in the isospin symmetric nuclear medium ($\eta = 0$) and in isospin asymmetric nuclear medium ($\eta = 0.5$), it changes to -29.2 MeV. The increase in the magnitude of the mass-shift, with density ρ_B , is because of the larger drop in the dilaton field χ at higher densities. However, with increase in the isospin asymmetry of the medium the magnitude of the mass-shift decreases because the drop in the dilaton field χ is less at a higher value of the isospin asymmetry parameter η . At the nuclear matter saturation density $\rho_B = \rho_0$, and for temperature $T = 0$, the mass-shift for $\psi(3686)$ is observed to be about -117 and -114 MeV for values of the $\eta = 0$ and 0.5 respectively, and for $\psi(3770)$, the values of the mass-shift are seen to be about -155 MeV and -150 MeV respectively. At $\rho_B = 4\rho_0$ and zero temperature, the values of the mass-shift for $\psi(3686)$ are modified to -436 MeV and -396 MeV for $\eta = 0$ and 0.5 respectively, and, for $\psi(3770)$, the drop in the masses are about -577 MeV and -523 MeV respectively. As mentioned earlier, the drop in the dilaton field, χ , at finite temperature is less than at zero temperature and this behaviour is reflected in the smaller mass-shifts of the charmonium states at finite temperatures as compared to the zero temperature case. At nuclear matter saturation density $\rho_B = \rho_0$, and for temperature $T = 100$ MeV, the values of the the mass-shift for the J/ψ meson are observed to be about -6.77 MeV and -6.81 MeV for isospin symmetric ($\eta = 0$) and isospin asymmetric ($\eta = 0.5$) nuclear medium respectively. At baryon density $\rho_B = 4\rho_0$, temperature $T = 100$ MeV, the mass-shift for J/ψ is observed to be -28.4 MeV and -27.2 MeV for isospin symmetric ($\eta = 0$) and isospin asymmetric ($\eta = 0.5$) nuclear medium respectively. For the excited charmonium states $\psi(3686)$ and $\psi(3770)$, the mass-shifts at nuclear matter saturation density $\rho_B = \rho_0$ and temperature $T = 100$ MeV, are observed to be -91.8 MeV and -121.4 MeV respectively, for isospin symmetric nuclear matter ($\eta = 0$) and -92.4 MeV and -122 MeV for the isospin asymmetric nuclear medium with $\eta = 0.5$. For baryon density $\rho_B = 4\rho_0$, and temperature $T = 100$ MeV, mass-shift for the charmonium states $\psi(3686)$ and $\psi(3770)$ are modified to about -386 MeV and -510 MeV respectively, for isospin symmetric ($\eta = 0$) and -369 MeV and -488 MeV for isospin asymmetric nuclear medium with $\eta = 0.5$.

For temperature $T = 150$ MeV and at the nuclear matter saturation density $\rho_B = \rho_0$,

the mass-shifts for the charmonium states $J/\psi, \psi(3686)$ and $\psi(3770)$ are seen to be -6.25 , -85 and -112 MeV respectively in the isospin symmetric nuclear medium ($\eta = 0$). These values are modified to -7.2 , -98 and -129 MeV respectively, in the isospin asymmetric nuclear medium with $\eta = 0.5$. At a baryon density $\rho_B = 4\rho_0$, the values of the mass-shift for $J/\psi, \psi(3686)$ and $\psi(3770)$ are observed to be -26.4 , -358 and -473 MeV in isospin symmetric nuclear medium ($\eta = 0$) and in isospin asymmetric nuclear medium with $\eta = 0.5$, these values are modified to -27.6 , -375 and -494 MeV respectively. Note that at high temperatures e.g at $T = 150$ MeV the mass-shift in isospin asymmetric nuclear medium ($\eta = 0.5$) is more as compared to isospin symmetric nuclear medium ($\eta = 0$). This is opposite to what is observed for the zero temperature case. The reason is that at high temperatures the dilaton field χ has more drop in the isospin asymmetric nuclear medium ($\eta = 0.5$) as compared to the isospin symmetric nuclear medium ($\eta = 0$), due to the contributions from the δ field for nonzero η , which is observed to decrease in its magnitude at high temperatures.

The values of the mass-shift for the charmonium states obtained within the present investigation, at nuclear saturation density ρ_0 and temperature $T = 0$, are in good agreement with the the mass shifts of J/Ψ , $\Psi(3686)$ and $\Psi(3770)$ as -8 , -100 and -140 MeV respectively, at the nuclear matter saturation density, computed in Ref. [8] from the second order QCD Stark effect, with the gluon condensate in the nuclear medium computed in the linear density approximation. The mass-shift for J/ψ has also been studied with the QCD sum rules in [6] and the value at nuclear saturation density was observed to be about -7 MeV. In Ref. [3] the operator product expansion was carried out upto dimension six and the mass shift for J/ψ was calculated to be -4 MeV at nuclear matter saturation density ρ_0 and at zero temperature. The effect of temperature on the J/ψ in the deconfinement phase was studied in [27, 32]. In these investigations, it was reported that J/ψ mass is essentially constant in a wide range of temperatures and above a particular value of the temperature, T , there was observed to be a sharp change in the mass of J/ψ in the deconfined phase [33]. In the present work, we have studied the effects of temperature, density and isospin asymmetry, on the mass modifications of the charmonium states ($J/\psi, \psi(3686)$ and $\psi(3770)$) in the confined hadronic phase, arising due to modifications of a scalar dilaton field which

simulates the gluon condensates of QCD, within a chiral SU(3) model. The effect of temperature was found to be small for the charmonium states $J/\psi(3097)$, $\psi(3686)$ and $\psi(3770)$, whereas, the masses of charmonium states observed to vary considerably with density, in the present investigation.

In summary, in the present work, we have investigated the effects of density, temperature and isospin asymmetry of the nuclear medium on the masses of the charmonium states J/ψ , $\psi(3686)$ and $\psi(3770)$, arising due to modification of the scalar dilaton field, χ , which simulates the gluon condensates of QCD, within the chiral SU(3) model and second order QCD Stark effect. The change in the mass of J/ψ with density is observed to be small at nuclear matter saturation density and is in agreement with the QCD sum rule calculations. There is seen to be an appreciable drop in the in-medium masses of excited charmonium states $\psi(3686)$ and $\psi(3770)$ with the density. At the nuclear matter saturation density, the mass shifts of these states are similar to the values obtained using the QCD second order Stark effect with the modifications of the gluon condensates computed in the linear density approximation [8]. For a given value of density and temperature, the effect of the isospin asymmetry of the medium on the in-medium masses of the charmonium states is found to be small. This is due to the fact that the magnitude of the δ field remains small as compared to the σ and ζ fields. At finite densities, the effect of the temperature on the charmonium states is found to decrease the values of mass-shift upto a particular temperature, above which the mass shift is seen to rise. This is because of an initial increase in the dilaton field χ and then a drop with further increase in the temperature, at a given baryon density, arising from solving the coupled equations for the scalar fields. This is related to the fact that the scalar densities of the nucleons initially drop and then rise with the temperature at finite values of the baryon densities. The mass drop of the excited charmonium states ($\Psi(3686)$ and $\Psi(3770)$) are large enough to be seen in the dilepton spectra emitted from their decays in experiments involving $\bar{p}$ -A annihilation in the future facility at GSI, provided these states decay inside the nucleus. The life time of the J/Ψ has been shown to be almost constant in the nuclear medium, whereas for these excited charmonium states, the lifetimes are shown to reduce to less than 5 fm/c, due to appreciable increase in their decay widths [29]. Hence a significant fraction of the produced excited charmonium states in these experiments are expected to decay inside

the nucleus [34]. The in-medium properties of the excited charmonium states $\psi(3686)$ and $\psi(3770)$ can be studied in the dilepton spectra in $\bar{p}$ -A experiments in the future facility of the FAIR, GSI. The mass shifts of the charmonium states in the hot nuclear medium seem to be appreciable at high densities as compared to the temperature effects on these masses, and these should show in observables like the production of these charmonium states, as well as of the open charmed mesons in the compressed baryonic matter (CBM) experiment at the future facility at GSI, where baryonic matter at high densities and moderate temperatures will be produced.

Acknowledgments

One of the authors (AM) is grateful to the Frankfurt Institute for Advanced Research (FIAS), University of Frankfurt, for warm hospitality and acknowledges financial support from Alexander von Humboldt Stiftung when this work was initiated. Financial support from Department of Science and Technology, Government of India (project no. SR/S2/HEP-21/2006) is also gratefully acknowledged.

-
- [1] J. P. Blaizot and J. Y. Ollitrault, Phys. Rev. Lett. **77**, 1703 (1996).
 - [2] T. Matsui and H. Satz, Phys. Lett. B **178**, 416 (1986).
 - [3] Sugsik Kim, Su Houn Lee, Nucl. Phys. A **679**, 517 (2001).
 - [4] Arata Hayashigaki, Phys. Lett. B **487**, 96 (2000).
 - [5] T. Hatsuda, S.H. Lee, Phys. Rev. C **46**, (1992) R34.
 - [6] F. Klingl *et al.*, Phys. Rev. Lett. **82**, 3396 (1999).
 - [7] M.E. Peskin, Nucl. Phys. **B156**, 365 (1979).
 - [8] Su Houn Lee and Che Ming Ko, Phys. Rev. C **67**, 038202 (2003).
 - [9] P. Papazoglou, D. Zschesche, S. Schramm, J. Schaffner-Bielich, H. Stöcker, and W. Greiner, Phys. Rev. C **59**, 411 (1999).
 - [10] A. Mishra, E. L. Bratkovskaya, J. Schaffner-Bielich, S. Schramm and H. Stöcker, Phys. Rev. C **70**, 044904 (2004). A. Mishra and S. Schramm, Phys. Rev. C **74**, 064904 (2006).

- [11] Amruta Mishra, Arvind Kumar, Sambuddha Sanyal, S. Schramm, Eur. Phys. J. A **41**, 205 (2009).
- [12] Amruta Mishra and Arindam Mazumdar, Phys. Rev. C **79**, 024908 (2009).
- [13] A. Mishra, E. L. Bratkovskaya, J. Schaffner-Bielich, S.Schramm and H. Stöcker, Phys. Rev. C **69**, 015202 (2004).
- [14] Arvind Kumar and Amruta Mishra, arXiv : 0912.2477v1 [nucl-th].
- [15] D. Zschesche, A. Mishra, S. Schramm, H. Stöcker and W. Greiner, Phys. Rev. C **70**, 045202 (2004).
- [16] A. Mishra, K. Balazs, D. Zschesche, S. Schramm, H. Stöcker, and W. Greiner, Phys. Rev. C **69**, 024903 (2004).
- [17] S.Weinberg, Phys. Rev. **166** 1568 (1968).
- [18] S. Coleman, J. Wess, B. Zumino, Phys. Rev. **177** 2239 (1969); C.G. Callan, S. Coleman, J. Wess, B. Zumino, Phys. Rev. **177** 2247 (1969).
- [19] W. A. Bardeen and B. W. Lee, Phys. Rev. **177** 2389 (1969).
- [20] J. J. Sakurai, Currents and Mesons, University of Chicago Press, Chicago, 1969.
- [21] O. Dumbrajs, R. Koch, L. Pilkuhn, G. C. Oades, H. Behrens, J. J. De Swart and P. Kroll, Nucl. Phys. **B 216**, 277 (1983).
- [22] Th. A. Rijken, V. G. J. Stoks, Y. Yamamoto, Phys. Rev. C **59**, 21 (1999); V. G. J. Stoks and Th. A. Rijken, Phys. Rev. C **59**, 3009 (1999).
- [23] G. Höhler, E. Pietarinen, I. Sabba-Stefanescu, F. Borkowski, G. G. Simon, V. H. Walther, R. D. Wendling, Nucl. Phys. **B 114**, 505 (1976).
- [24] J. Schechter, Phys. Rev. D **21**, 3393 (1980).
- [25] Erik K. Heide, Serge Rudaz and Paul J. Ellis, Nucl. Phys. A **571**, (2001) 713.
- [26] Thomas D. Cohen, R. J. Furnstahl and David K. Griegel, Phys. Rev. C **45**, 1881 (1992).
- [27] Su Houn Lee and Kenji Morita, Phys. Rev. D **79**, 011501(R) (2009).
- [28] Xuemin Jin, Thomas D. Cohen, R. J. Furnstahl and David K. Griegel, Phys. Rev. C **45**, 1881 (1992).
- [29] Bengt Friman, Su Houn Lee and Taesoo Song Phys. Lett. **B 548**, 153 (2002).
- [30] R. J. Furnstahl and B. D. Serot, Phys. Rev. C **41**, 262 (1992); J. Theis, G. Graebner, G.

- Buchwald, J. Maruhn, W. Greiner and J. Polonyi, Phys. Rev. D **28**, 2286 (1983).
- [31] G.Q.Li, C. M. Ko and G. E. Brown, Nucl. Phys. **A 606**, 568 (1998).
- [32] Cesareo A. Dominguez, Marcelo Loewe, J. Cristobal Rojas and Yingwen Zhang arXiv 0908.2709 [hep-ph].
- [33] Kenji Morita and Su Houng Lee, Phys. Rev. Lett. **100**, 022301 (2008).
- [34] Y. S. Golubeva et al, nucl-th/0212074.